

Please remember BLINCYTO may cause:

1. Neurological events, including immune effector cell-associated neurotoxicity syndrome (ICANS). Signs and symptoms of this side effect may include:
 - » Uncontrolled shaking (tremor)
 - » Confusion
 - » Disturbances of brain function (encephalopathy)
 - » Difficulty in communicating (aphasia) and/or writing (dysgraphia)
 - » Seizure (convulsion)
2. Cytokine release syndrome, including:
 - » Fever
 - » Chills
 - » Swelling
 - » Dizziness or lightheadedness



Please seek urgent attention from your doctor or nurse immediately or seek emergency help if you experience any of these symptoms



Please travel home safely and do not drive or operate moving vehicles/heavy machinery, or engage in hazardous activities whilst receiving BLINCYTO

Patient Card



Please show this card to all emergency and healthcare providers

**Information about
BLINCYTO®▼(blinatumomab)**

My name is _____

In Case of Emergency (ICE) contact _____

I am being treated with BLINCYTO, a treatment for B-precursor acute lymphoblastic leukaemia, which can lower my immune system.

I started treatment on _____

Before providing any treatment, please call my prescribing physician at the number below. If any medical evaluations are undertaken, please provide copies of all medical records, including any treatments and/or test results, to the doctor(s) named below.

	Name	Hospital	City	Phone Number
Haematologist				
Oncologist				
Haematology Nurse				

▼This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.