






<p>This Alert Card contains important safety information that you need to be aware of before and during treatment with Flixabi.</p> <p>Please read the Flixabi ‘Package Leaflet’ carefully before you start using this medicine.</p> <p>It is important that you and your doctor record the brand name and batch number of your medicines.</p>	<p><b>Infections</b></p> <p><b>Before treatment with Flixabi</b></p> <ul style="list-style-type: none"> <li>• Tell your doctor if you have an infection even if it is a very minor one.</li> <li>• It is very important that you tell your doctor if you have ever had tuberculosis (TB), or if you have been in close contact with someone who has had TB. Your doctor will test you to see if you have TB. Ask your doctor to record the type and date of your last screening(s) for TB on the card.</li> </ul>	<ul style="list-style-type: none"> <li>• Tell your doctor if you have hepatitis B or if you know or suspect you are a carrier of the hepatitis B virus.</li> </ul> <p><b>During treatment with Flixabi</b></p> <ul style="list-style-type: none"> <li>• Tell your doctor straight away if you have signs of an infection. Signs include a fever, feeling tired, (persistent) cough, shortness of breath, weight loss, night sweats, diarrhea, wounds, dental problems, burning when urinating or ‘flu-like’ signs.</li> </ul>	<p><b>Heart Failure</b></p> <p><b>Before treatment with Flixabi</b></p> <ul style="list-style-type: none"> <li>• Tell your doctor if you have any heart problems such as mild heart failure.</li> </ul> <p><b>During treatment with Flixabi</b></p> <ul style="list-style-type: none"> <li>• Tell your doctor straight away if you notice signs of a heart problem. Signs include shortness of breath, swelling of the feet or changes in your heartbeat.</li> </ul>	<p><b>Pregnancy and Vaccinations</b></p> <ul style="list-style-type: none"> <li>• In case you have received Flixabi while you were pregnant, it is important that you inform your baby’s doctor about it before your baby receives any vaccine. Your baby should not receive a ‘live vaccine’, such as BCG (used to prevent tuberculosis) within 6 months after birth.</li> </ul>
--	---	---	--	---

PROFILE NAME		PRODUCT(S)		PATTERNS AND COLORS	
PATIENT ALERT CARD_1 LANG_SB2_V01_P5		SB2		 TRANSPARENT LABEL	 PROFILE  TEXT LIMIT
DIMENSION mm		DATE MODIFIED			
85 x 226mm		25 04 2016			
SUPPLIER DIE LINE		PREPARED BY:			
SCHAWKI		SCHAWK			 PROFILE (Non Print)  Technical (Non Print)

ITEM NUMBER	DESCRIPTION			PRIMARY FONT USED		ARTWORK COLORS					
48845-01	ALLIANCE PATIENT ALERT CARD SB2 CONC 100MG GBR/IRL			Calibri, Calibri-Bold		<div><div><div>Black</div><div>PMS 166</div><div>PMS 286</div><div>PMS 375</div><div>Technical Ink</div></div><div><div><div>White</div><div>Varnish</div></div></div></div> <div>biogen</div>					
	LANGUAGE(S)		TEMPLATE NUMBER	MIN FONT SIZE (BODY TEXT)							
	English		N/A	6.0 pt							
PROOF NUMBER	PHARMA CODE	PHARMA CODE TYPE	VERSION CODE	GS REFERENCE	DATE MODIFIED						
04	1264	Standard	N/A	Schawk-517294A05	27 April 2016						
	BARCODE NUMBER		CMO ITEM NUMBER	APPROVED BY							
	N/A		255952								

